

NIRVIN

Drops & Spray

Composition

Nirvin adult drops

Each ml contains Xylometazoline HCl 1.0 mg

Nirvin children drops

Each ml contains Xylometazoline HCl 0.5 mg

Nirvin adult spray

Each ml contains Xylometazoline HCl 1.0 mg

Nirvin children spray

Each ml contains Xylometazoline HCl 0.5 mg

Action & Indication

It is a sympathomimetic with marked alphasadrenergic action. It acts as a vasoconstrictor when applied topically to mucous membrane & thus reduces swelling & congestion. Effect begins within 0 min of application & lasts for up to 10 hrs. It is a useful nasal decongestant &, in lower strength a good conjunctival decongestant.

Warnings

If you are using this medicine without a prescription, carefully read and follow any precautions on the label. For Xylometazoline, the following should be considered:

Allergies

Tell your doctor if you have ever had any unusual or allergic reaction to Xylometazoline or to any of the other nasal decongestants. Also, tell your health care professional if you are allergic to any other substances, such as foods, preservatives, or dyes.

Children

Children may be especially sensitive to the effects of Xylometazoline. This may increase the chance of side effects during treatment, especially sedation.

Adults

Many medicines have not been studied specifically in older people. Therefore, it may not be known whether they work exactly the same way they do in younger adults or if they cause different side effects or problems in older people. There is no specific information comparing use of Xylometazoline in the elderly with use in other age groups.

Other medical problems

The presence of other medical problems may affect the use of Xylometazoline. Make sure you tell your doctor if you have any other medical problems, especially:

Diabetes mellitus (sugar diabetes)—Xylometazoline may worsen circulatory problems.

Enlarged prostate—Urination difficulty may worsen due to constrictive effects of Xylometazoline.

Glaucoma—Xylometazoline may worsen the circulatory problems.

Heart or blood vessel disease or high blood pressure—Xylometazoline may make the condition worse
Overactive thyroid

Adverse Reactions

Along with its needed effects, a medicine may cause some unwanted effects. Although not all of these side effects may occur, if they do occur they may need medical attention.

When this medicine is used for short periods at low doses, side effects usually are rare. However, check with your doctor as soon as possible if any of the following occur:

Increase in runny or stuffy nose

Symptoms of too much medicine being absorbed into the body

Blurred vision; dizziness; headache or light-headedness; hives ; nausea; nervousness; pounding, irregular, or fast heartbeat; rash; shortness of breath; swelling of eyelids, face, or lips; troubled breathing ; trouble in sleeping; wheezing

Other side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. However, check with your health care professional if any of the following side effects continue or are bothersome:

Burning, dryness, or stinging of inside of nose; discharge of fluid from the nose; sneezing

Proper Use of This Medicine

To use the nose drops

Blow your nose gently. Tilt the head back while standing or sitting up. Place the drops into each nostril and immediately bend head forward toward the knees for a few seconds to allow the medicine to spread throughout the nose.

Rinse the dropper with hot water and dry with a clean tissue. Replace the cap right after use.

To use the nose spray

Blow your nose gently. With the head upright, spray the medicine into each nostril. Sniff briskly while squeezing the bottle quickly and firmly. For best results, spray once into each nostril, wait 3 to 5 minutes to allow the medicine to work, and then blow your nose gently and thoroughly. Repeat until the complete dose is used.

Rinse the tip of the spray bottle with hot water taking care not to suck water into the bottle, and dry with a clean tissue. Replace the cap right after use.

To avoid spreading the infection, do not use the container for more than one person. Use this medicine only as directed

Dosage

The dose of nasal Nirvin will be different for different patients. Follow your doctor's orders or the directions on the label. The following information includes only the average doses of nasal Nirvin. If your dose is different, do not change it unless your doctor tells you to do so.

For stuffy nose

For nasal solution (nose drops or nose spray) dosage forms:

Adults and children 12 years of age and older

Use one to three drops or sprays of a 0.1% solution in each nostril every eight to ten hours as needed.

Children 2 to 12 years of age

Use two or three drops or sprays of a 0.05% solution in each nostril every eight to ten hours as needed.

Children up to 2 years of age

Use and dose must be determined by your doctor.

Missed dose

If you miss a dose of this medicine and you remember within an hour or so of the missed dose, use it right away. However, if you do not remember until later, skip the missed dose and go back to your regular dosing schedule. Do not double doses.

Storage

Keep out of the reach of children. Store away from heat and direct light.

Keep the medicine from freezing. Do not keep outdated medicine or medicine no longer needed. Be sure that any discarded medicine is out of the reach of children.

Presentation

Nirvin adult drops

Bottle of 10 ml

Nirvin children drops

Bottle of 10 ml

Nirvin adult spray

Metered dosing bottle of 10 ml

Nirvin children spray

Metered dosing bottle of 10 ml