

TETRIN

Eye Drops

Composition

Tetrin Eye Drops

Each ml contains Tetrahydrozoline HCl 0.05%.

Action

Tetrin Eye Drops is a sterile, stable, non-irritating solution. Its sympathomimetic action promotes decongestion.

Indications

For the relief of burning, itching, irritation and redness of eyes.

Contraindications

- Should not be used by individuals with glaucoma or other serious eye disease.
- It should not be used with monoamine oxidase inhibitors or within 10 days of their discontinuation, since hypertensive crisis may result.

Adverse Reactions

Some side effects may appear during treatment with this drug such as: Pain in Eyes, change in eye sight, irritation and redness of eyes.

Warnings and Precautions

Tetrahydrozoline should be used only in minor eye irritation. If relief is not obtained within 48 hours, or irritation persists or increases, discontinue use. Infection, pus, the presence of a foreign body in the eye and any mechanical, chemical, heat or other injury to the eye requires the immediate attention of the physician.

As the possibility of adverse effects on the corneal permeability and the danger of disruption of the corneal epithelium with prolonged or repeated usage of benzalkonium chloride preserved ophthalmological preparations cannot be excluded, regular ophthalmological examination is required. Caution should be exercised in the use of benzalkonium chloride preserved topical medication over an extended period in patients with extensive ocular surface disease.

Do not let the applicator tip touch the eye or any other surface, since this may contaminate the eye drops. Tetrahydrozoline must be used with caution in patients with hypertension and hyperthyroidism.

Pregnancy

Category A

Adequate and well-controlled studies have failed to demonstrate a risk to the fetus in the first trimester of pregnancy (and there is no evidence of risk in later trimesters).

Tetrahydrozoline is safe during pregnancy.

Nursing Mothers

Tetrahydrozoline is safe for use by nursing mothers

Drug Interactions

There are no known drug interactions with Tetrahydrozoline.

Dosage and Administration

Instil one or two drops of Tetrin Eye Drops in each eye two or three times a day. Tetrin Eye Drops should be used sparingly in children. Its use in children under 2 years of age is not recommended.

DIRECTIONS FOR USE

Shake well before use. Wash hands carefully. Pull the head backwards or lie on a bed and with the aid of a finger, pull the lower eyelid down. Drip the medicine into the space formed and close your eyes, for 1-2 minutes. Do not blink. If you feel that a drop has not been properly absorbed, add an additional drop. Wipe the remaining liquid around the eye with a tissue. To avoid contamination of the solution pay attention that the tip of the bottle does not come in contact with any Surface, including the finger or eye, and Close the bottle tightly.

Storage Conditions

Store in a cool place below 25°C and protect from light. Discard any remaining medicine after 30 days of use.

Presentation

Tetrin Eye drops

Bottle of 10 ml