

# PANTINOL PLUS

Cream

## Composition

Each 1 gram contains 50 mg Dexpanthenol and 5 mg Chlorhexidine HCl

## Action

Pantinel plus cream prevents and fights infections, while promoting healing of the skin. Is applied on the skin and absorbed easily from skin. It doesn't leave stain to the clothes.

## Mechanism of Action

Dexpanthenol, the active ingredient of Pantinel plus Cream is a precursor Pantothenic acid, one of B complex vitamins. It is absorbed easily on the skin and converted to Pantothenic acid.

Pantothenic acid is essential for skin and mucosa. It helps skin regeneration, heal skin rashing and irritation and accelerate the healing of minor wounds.

Chlorhexidine HCl is a disinfectant and plus against to the most of bacteria and some fungus.

It is especially effective to Staphylococcus aureus which causes skin infections.

## Indications

- In case of minor injuries of any kind, that could become infected: scrapes, cuts, scratches, cracks, burns.
- Light, intertrigo (inflammation of the skin at the folds).
- In case of inflammation of the skin accompanied by an infection;
- In case of inflammation of the nipple during lactation;
- On prescription only in cases of chronic wounds such as leg ulceration (ulcus cruris), decubitus ulcer and surgical wounds.

## Contraindications

- It shouldn't be used in patients with known hypersensitivity to any components
- Injury to the eardrum, Pantinel Plus should not be applied in the ear.
- In the presence of acute wounds, deep and very dirty, such injuries must be treated by the doctor.

## Adverse Reactions

In rare case allergic reactions can occur.

## Warnings and Precautions

It must not be applied to the eyes and mucosa. In the presence of extensive wounds, deep and very dirty, such as those caused by bites, or pointed objects, intervention by the physician is required (risk of tetanus). If the size of the wound remains the same for some time or if the healing does not occur in the space of 10-14 days, it is also necessary to consult a doctor. This applies even if the edges of the wound are very red, or if the wound suddenly starts to swell, so it is very painful or if the injury is accompanied by fever (risk of blood poisoning).

## Pregnancy and lactation

Pantinel Plus can be used during pregnancy and lactation. However, do not apply over large areas during this period.

## Dosage and Administration

Gently rub the medicine in until it is evenly distributed to the affected area.

- **Wound and burns:** 1 time daily or if necessary more times.
- **Nipple cracked in nursing mothers:** After every breastfeed applied to nipples. Before nursing, nipples must be cleaned with water.
- **Nappy rash:** Pantinel plus cream is used every nappy change.

**Storage**

Keep tube tightly closed and store at room temperature below 30° C. Keep away from children.

**Presentation**

Tube of 25 grams